

Performance Improvement Summary

| Focus | Target | January | February | March | April | May | June | July | Aug | Sept | Oct | Nov | Dec |
|---|---------------|---------|----------|-------------|--------|----------------------|--------|------|----------------------|--------|-----|-----|-----|
| HTN | 68% | | | 66.71% | | | 71.20% | | | | | | |
| <i>Percentage of patients with diagnosed hypertension whose most recent blood pressure was less than or equal to 140/90</i> | | | | | | | | | | | | | |
| Cervical Cancer | 60% | 12.55% | 12.00% | | 42.14% | | | | 36.04% | | | | |
| <i>Percentage of women 21-64 years of age who received one or more tests to screen for cervical cancer</i> | | | | | | | | | | | | | |
| HGA1C <9% | 70% | | | 56.77% | | | 44.71% | | | | | | |
| <i>Percentage of diabetic patients with HBA1C levels less than or equal to 9%</i> | | | | | | | | | | | | | |
| Immunization | 75% | | | | | 18.40% | | | | | | | |
| <i>Percentage of children with 2nd birthday during the measurement year with appropriate immunizations.</i> | | | | | | | | | | | | | |
| BHI | 60% | | | | | | | | | | | | |
| <i>Ratio of patients that depression screening indicated need for BHI and that received access to BHI services during a medical visit.</i> | | | | | | | | | | | | | |
| Oral Health | 10% | | | | | | | | | | | | |
| <i>Percentage of patients with a filling needing to be replaced or refilled within a year.</i> | | | | | | | | | | | | | |
| Dental Sealants | 28% | | | | | | | | | | | | |
| <i>Number of patients ages 6-9 at elevated risk for cavities who received sealants on permanent first molars</i> | | | | | | | | | | | | | |
| Asthma | 85.00% | 0% | | | | | | | 11.28% | | | | |
| <i>Percentage of patients age 5-40 years with a diagnosis of persistent asthma who were prescribed long term control medication during the current year.</i> | | | | | | | | | | | | | |
| CAD | 65% | | 57.00% | | 60.43% | | | | 58.90% | 58.77% | | | |
| <i>Percentage of patients 18+ with a diagnosis of CAD prescribed a lipid lowering therapy.</i> | | | | | | | | | | | | | |
| Adolescent Weight | 40.00% | 5.80% | | | | 98.42% 46.70% 10.30% | | | 98.89% 67.97% 26.78% | | | | |
| <i>Percentage of pts 2-17 years of age who had a visit with evidence of BMI percentile documentation, counseling for nutrition and physical activity.</i> | | | | | | | | | | | | | |
| IVD | 70% | | 17.25% | | 14.03% | | 18.54% | | | 23.33% | | | |
| <i>Patients 18+ with a diagnosis of Ischemic Vascular Disease (IVD) who had documentation of use of aspirin or antithrombotic during the measurement year.</i> | | | | | | | | | | | | | |
| Colorectal Cancer | 30% | | 30.00% | | 32.00% | | 30.72% | | | 30.39% | | | |
| <i>Percentage of patients ae 50-70 with appropriate screening for colorectal cancer including colonoscopy (10 yrs), sigmoidoscopy (5 yrs) or annual fecal testing.</i> | | | | | | | | | | | | | |
| Tobacco Cessation | 30% | 74% | | | 80% | | | | 79.65% | | | | |
| <i>Percentage of patients 18+ that have been seen for at least 2 office visits who have been queried about tobacco use one or more times within 24 mos and received cessation counseling.</i> | | | | | | | | | | | | | |
| Depression Screening | 60% | 68.50% | 69.00% | | 69.73% | 71.40% | | | 71.01% | | | | |
| <i>Percentage of patients ages 12+ that have been screened for behavioral health needs and a follow-up plan identified.</i> | | | | | | | | | | | | | |
| HIV | 75% | | | | | | | | | | | | |
| <i>Percentage of newly diagnosed HIV patients who had a medical visit for HIV care within 90 days of a diagnosis</i> | | | | | | | | | | | | | |
| Perinatal | 7.80% | | | | 38.00% | | | | | | | | |
| <i>Percentage of births less than 2,500 grams to health center patients</i> | | | | | | | | | | | | | |
| Prenatal | 77.90% | | | 100.00% | | | | | | | | | |
| <i>Percentage of women beginning prenatal care in the first trimester.</i> | | | | | | | | | | | | | |
| Adult Weight Measure | 26.00% | | | 74.34/77.41 | | | | | 83.68/79.92% | | | | |
| <i>% of pts age 18 + with (1) BMI charted and (2) follow up plan if pts are overweight or underweight, as documented in the HER</i> | | | | | | | | | | | | | |